

2. DESCRIPTIONS OF SEVEN NEW SPECIES OF BIRDS DISCOVERED
BY MR. J. J. MONTEIRO IN THE PROVINCE OF BENGUELA,
ANGOLA, WEST AFRICA. BY DR. G. HARTLAUB, FOREIGN
MEMB.

(Plates IV., V., VI.)

1. CRATEROPUS GYMINOGENYS, sp. nov.

Capitis lateribus circumscripte nudis; pileo albido, nucham versus magis cinerascente; dorso fuscō; collo postico et lateralē lātius rufescēte; alis et cauda nigro-fuscis; uropygio albo, fulvescente lavato; subalaribus dilute rufis; lateribus rufescētibus; subcaudalibus albīs; remigum marginib⁹ internis pallide rufis, primo et secundo exceptis totis nigris; rostro et pedibus nigris.
Long. circa 9", rostri a fr. $9\frac{1}{4}$ ", alae 3" 11", caudæ 3" 8", tarsi $14\frac{1}{2}$ ".

2. DRYOSCOPUS GUTTATUS, sp. nov.

Supra nitide niger, nitore chalybeo; cauda tota nigerrima; alis nigris, macula scapulari transversa alba; marginibus externis remigum duorum secundariorum late niveis, fasciam angustam longitudinalē formantibus; subalaribus albīs, flexura alae nigra; tergi et uropygii plumis obscure cinerascentibus, macula anteapicali rotundata alba notatis, ipso margine apicali nigro; abdomine unicolore albo; subcaudalibus albīs; rostro valde compresso, vix plus quam emarginatum, nigro; pedibus nigris.
Long. 7" 8", rostri a fr. 8", alae 3" 7", caudæ 3" 7", tarsi 15".
Obs. A typical species.

3. UPUPA DECORATA, sp. nov.


Simillima U. capensi, sed diversa; remigibus secundariis albīs ante apicem extensem nigrum, nigro bifasciatīs, fasciola nigredinis apicalis medio interrupta; abdomine in fundo pallide rufescēte, maculis nonnullis longitudinalibus nigro-fuscis.

Long. rostri a fr. 1" 7", alae 5", caudæ 4", tarsi 8".

Obs. The unique specimen of this bird seems to be not quite adult; the crest is exactly like that of *U. capensis*; the rufous colour of the under parts goes much further down in *U. capensis*. This latter species has the secondary remiges, from the base to the black apical part, of a pure white.

4. TOCCUS ELEGANS, sp. nov. (Pl. IV.)

Pileo ardesiaco, occipite subcristato; capitis lateribus albīdis, ardesiaco longitudinaliter variegatis, superciliis eodem modo pictis; dorso fuscescenti-nigro, area mediana angusta alba; alarum tectricibus in fundo nigro maculis magnis rotundatis albīs, nigerrime marginatis, pulchre variis; remigibus majoribus nigris, 1° et 2° macula alba unica in pogonio interno, 3°-7° albo bimaculatis, 3 secundariis albīs, macula magna irregulari



J. Wolf del. et lith.

M. & N. Hanhart imp.

TOCCUS ELEGANS.

